


BEST AVAILABLE COPY

		Minimum	Maximum	Ca Average	Cap color e standard dev.	3	Color indication
Broncoh	_	64.83	72.24	69.67	1.62	20	
	а	5.87	7.33	6.61	0.34	20	
	b	19.31	21.72	20.82	0.59	20	
C9	L	59.46	68.20	63.41	2.11	20	
	а	7.60	9.27	8.55	0.40	20	
	d	17.80	20.94	19.55	0.83	20	
U1	L	90.12	95.16	94.14	1.21	20	
	a	0.60	1.24	0.88	0.15	20	
	b	6.39	8.32	7.20	0.53	20	
				Stip	Stipe color		
		Minimum	Maximum	Average	standard dev.	n	Color indication
Broncoh	Г	84.10	89.45	86.93	1.67	20	
	а	0.06	1.72	0.96	0.48	20	
	Ь	7.31	10.49	8.84	0.98	.20	意思を不可能という。
ග	Г	82.87	90.56	87.58	1.73	20	
	а	0.09	1.83	0.57	0.54	20	
	σ	7.74	15.78	9.37	1.78	20	
U	_	83.09	90.23	86.66	2.62	20	
	മ	0.48	1.99	1.06	0.42	20	
	σ	6.44	10.75	7.91	1.10	20	

AVAILABLE COPY

Mushroom Variety Named 'Broncoh' Inventor: Anton Sonnenberg Attorney Docket No.: 2475-040292



BEST AVAILABLE COPY

Fig. 5

Mushroom Variety Named 'Broncoh' Inventor: Anton Sonnenberg Attorney Docket No.: 2475-040292






Fig. 7

BEST AVAILABLE COPY

		<u>_</u>			69			Bronc oh				<u>_</u>			C9			oh i	Bronc		Strain
σ	മ	_	Б	മ	_	Ь	മ	_		ь	a	Г	ь	മ	_	σ	B	_			
7.20	2.38	89.93	7.13	2.55	89.91	7.20	2.00	89.99		12.90	19.57	68.32	13.10	17.01	70.36	13.50	16.53	70.01		c	Duration of storage (hrs)
6.70	3.29	89.03	7.22	0.72	89.97	6.73	3.08	88.67	Discolo	15.03	18.82	65.27	18.26	3.99	72.20	15.23	16.99	65.53		Discolora	f storage s)
0.50	-0.92	0.89	-0.09	1.83	-0.06	0.48	-1.09	1.32	Discoloration of flesh	-2.13	0.75	3.05	-5.16	13.02	-1.83	-1.73	-0.46	4.49		2 Discoloration of lamellan	difference
· · · · · · · · · · · · · · · · · · ·			まるなどのなっていた。						sh											0	Color indication
			いてのでは、東京の地域では、アイン	***																N	dication

BEST AVAILABLE COPY